

Scattering database of oriented dust particles with realistic shapes and sizes

Alexandra Tsekeri^{a,*}, Josef Gasteiger^b, Alexander Konoshonkin^{c,d}, Natalia Kustova^c,
Thanasis Georgiou^a, and Vassilis Amiridis^a

^a*Institute for Astronomy, Astrophysics, Space Applications and Remote Sensing, National Observatory of Athens, Athens, Greece*

^b*Faculty of Physics, University of Vienna, Vienna, Austria*

^c*National Research Tomsk State University, Tomsk, Russia*

^d*V. E. Zuev Institute of Atmospheric Optics, SB RAS, Tomsk, Russia*

*Presenting author (atsekeri@noa.gr)

Dust orientation is an ongoing investigation in recent years [1]. Its potential proof will be a paradigm shift for dust remote sensing, invalidating the currently used simplifications of randomly-oriented particles. For the effective monitoring of the phenomenon we constructed a new scattering database that takes into account realistic-shaped oriented dust particles and a size range that includes most of the coarse mode of dust. In particular, we have used the irregular shapes of Gasteiger *et al.* [2], a size parameter range of 0.01 – 60 and four refractive indices with values of $1.48 - 1.6 + i0 - 0.002$. The scattering calculations were performed with the Amsterdam Discrete Dipole Approximation (ADDA) [3] using ~50M CPU core hours on High Performance Computing (HPC) systems.

Currently, there is no complete solution for calculating the scattering properties of the whole range of dust sizes, shapes and refractive indices. For size parameters exceeding 60 – 70, ADDA calculations are challenging even for HPC machines, due to their high computational cost [3]. For this reason we investigate the applicability of the much faster Physical Optics (PO) approximation [4] for dust particles with faceted shapes. We compare the PO scattering calculations with ADDA calculations and we present here the issues and perspectives of this coupling in formulating a complete scattering database for oriented dust.

References

- [1] Ulanowski, Z., J. Bailey, P. W. Lucas, *et al.*, 2007: Alignment of atmospheric mineral dust due to electric field. *Atmos. Chem. Phys.* **7**, 6161–6173.
- [2] Gasteiger, J., M. Wiegner, S. Gross, *et al.*, 2011: Modeling lidar-relevant optical properties of complex mineral dust aerosols. *Tellus B* **63**, 725–741.
- [3] Yurkin, M. A., and A. G. Hoekstra, 2011: The discrete-dipole-approximation code ADDA: capabilities and known limitations. *J. Quant. Spectrosc. Radiat. Transf.* **112**, 2234–2247.
- [4] Konoshonkin, A., A. Borovoi, N. Kustova, *et al.*, 2017: Light scattering by ice crystals of cirrus clouds: from exact numerical methods to physical-optics approximation. *J. Quant. Spectrosc. Radiat. Transf.* **195**, 132–140.

Preferred mode of presentation: Oral/Poster